

# IOWA STATE UNIVERSITY

## Digital Repository

---

Leopold Center Completed Grant Reports

Leopold Center for Sustainable Agriculture

---

2008

## Integration of natural seed treatments in organic and open-pollinated corn systems

A. Susana Goggi

*Iowa State University*, [susana@iastate.edu](mailto:susana@iastate.edu)

Kathleen Delate

*Iowa State University*, [kdelate@iastate.edu](mailto:kdelate@iastate.edu)

Linda M. Pollak

*United States Department of Agriculture*

Kendall R. Lamkey

*Iowa State University*, [krlamkey@iastate.edu](mailto:krlamkey@iastate.edu)

Follow this and additional works at: [http://lib.dr.iastate.edu/leopold\\_grantreports](http://lib.dr.iastate.edu/leopold_grantreports)



Part of the [Agronomy and Crop Sciences Commons](#), [Entomology Commons](#), [Horticulture Commons](#), and the [Plant Pathology Commons](#)

---

### Recommended Citation

Goggi, A. Susana; Delate, Kathleen; Pollak, Linda M.; and Lamkey, Kendall R., "Integration of natural seed treatments in organic and open-pollinated corn systems" (2008). *Leopold Center Completed Grant Reports*. 295.

[http://lib.dr.iastate.edu/leopold\\_grantreports/295](http://lib.dr.iastate.edu/leopold_grantreports/295)

This Article is brought to you for free and open access by the Leopold Center for Sustainable Agriculture at Iowa State University Digital Repository. It has been accepted for inclusion in Leopold Center Completed Grant Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Integration of natural seed treatments in organic and open-pollinated corn systems

## **Abstract**

Essential oils from the aromatic plants, such as thyme, oregano, cinnamon, clove, and savory, were used to effectively control several common seed and soil pathogens in the laboratory, and could be used as seed treatments in corn. Results from field experiments showed some oils protected the seeds from pathogens but not others. Before these natural products can be widely adopted, however, issues of volatility of the oils at ambient temperature and appropriate application rate must be addressed.

## **Keywords**

Agronomy, Horticulture, Biocontrol and Integrated Pest Management, Organic production practices and comparisons

## **Disciplines**

Agronomy and Crop Sciences | Entomology | Horticulture | Plant Pathology



LEOPOLD CENTER  
FOR SUSTAINABLE AGRICULTURE

# Integration of natural seed treatments in organic and open-pollinated corn systems

**Abstract:** Essential oils from the aromatic plants, such as thyme, oregano, cinnamon, clove, and savory, were used to effectively control several common seed and soil pathogens in the laboratory, and could be used as seed treatments in corn. Results from field experiments showed some oils protected the seeds from pathogens but not others. Before these natural products can be widely adopted, however, issues of volatility of the oils at ambient temperature and appropriate application rate must be addressed.

## Question & Answer

**Q:** What are the options for organic farmers who are interested in organic seed treatments?

**A:** This work shows that while the essential oils of thyme, oregano, cinnamon, clove, and savory don't harm seeds and effectively control three pathogens in the lab, it is still not proven that they would be effective in the field.

## Background

The number of alternative crop production systems has increased in recent years in response to growing concerns about agricultural concentration and interest in a more ecological, farm-based agriculture. In these low-input systems, planting dates for open-pollinated corn may be delayed to avoid the cold, wet spring soils. Conventional farmers use synthetic fungicide treatments to protect the emerging seedlings, but there are few organic fungicides available. The objective of this project was to determine whether the essential oils of aromatic plants can be used successfully as a contact fungicide seed treatment for organic corn.

## Approach and methods

Eighteen plant essential oils were screened for their

ability to control three common corn pathogens: *Penicillium* ear rot, *Fusarium* ear and root rot, and *Pythium* seedling blight and stalk rot. The oils screened were: anise (*Pimpinella anisum*), basil (*Ocimum basilicum*), bay (*Laurus nobilis*), caraway (*Carum carvi*), cinnamon (*Cinnamomum zeylanicum*), clove (*Eugenia caryophyllata*), coriander (*Coriandrum sativum*), lavender (*Lavendula officinalis*), lemongrass (*Cymbopogon citratus*), nutmeg (*Myristica fragrans*), oregano (*Origanum minutiflorum*), pepper (*Piper nigrum*), peppermint (*Mentha piperita*), rosemary (*Rosmarinus officinalis*), sage (*Salvia officinalis*), savory (*Satureia montana*), tea tree (*Melaleuca alternifolia*), and thyme (*Thymus vulgaris*). An initial concentration of 1,000 ppm of the essential oils was used to determine their effect on pathogen growth in the laboratory.

To determine the minimum inhibitory concentration (MIC, lowest concentration for pathogen control), the oils were diluted to concentrations of 100, 200, 400, and 800 ppm. To determine the presence of phytotoxicity problems related to use of the oils, artificially infected corn seeds were treated with rates up to 16,000 ppm (MIC x 20) of the essential oils and germinated.

## Results and discussion

Of the initial 18 plant essential oils screened, five (cinnamon, clove, oregano, savory and thyme) completely inhibited the growth of the three corn pathogens (*Fusarium* sp., *Penicillium* sp., and *Pythium* sp.) in vitro. The minimum

### Principal Investigator:

Susana Goggi  
Agronomy

### Co-investigators:

Kathleen Delate  
Agronomy and Horticulture  
Iowa State University

Linda Pollak

USDA-Agricultural Research  
Service

Ames

Kendall Lamkey

Agronomy

Iowa State University

### Budget:

\$25,000 for year one  
\$25,000 for year two

inhibitory concentration of the essential oils in the laboratory was 800 ppm. The growing seedlings were not affected (i.e., no phytotoxicity symptoms were seen) at rates up to 16,000 ppm concentration of the essential oils.

Gas chromatography analysis determined the major constituent compounds of the five oils. The antifungal properties of the essential oils seemed to be associated with aromatic compounds. The concentration of these compounds varied among the oils and from those reported in the literature. Further research should be conducted comparing the efficacy of the pure forms of the constituent compounds and the essential oil as a whole.

Field emergence of inbred and hybrid seed treated with essential oils was significantly lower than seed treated with commercial fungicides Maxim XL, a conventional fungicide, and Natural 2, an organic fungicide; however, the emergence rates were not different from the organic fungicide Yield Shield or an untreated control.

The standard germination test was used to evaluate the germination of artificially infected inbred and hybrid corn seeds treated with two rates of the five essential oils. There

were no significant differences in germination percentage between the two rates of essential oil treatments. The germination percentage of seeds treated with the essential oil was significantly lower than the commercial seed treatments Natural 2 and Maxim XL. However, some essential oil seed treatments were not significantly different from the commercial organic seed treatment Yield Shield.

## Conclusions

The essential oils of cinnamon, clove, oregano, savory, and thyme control common seed and soil borne pathogens in the laboratory at very low (800 ppm) rates; are not phytotoxic to the growing seedlings; and can potentially be used as seed treatments. In the laboratory and field experiments, the essential oils are not as effective, possibly due to volatilization and consequent loss of the essential oils. Further research should be conducted to determine the correct application rate for the essential oils and possible methods to reduce the volatilization of the aromatic compounds from the seed surface.


## Impact of results

The essential oils of thyme, oregano, cinnamon, clove, and savory effectively controlled three pathogens *in vitro*. *In vivo* results were mixed and further research is necessary to determine the true potential of these essential oils as seed treatments.

## Education and outreach

Results from these experiments were shared with farmers and colleagues at Practical Farmers of Iowa and the Michael Fields Institute at their annual farmer meetings and field demonstrations. Results also were reported at the ISU Seed Science Center-sponsored workshops, including the seed quality workshops.

An article based on the project findings on aromatic plant oils as a fungicide for organic corn production was to appear in *Crop Science* journal. (See: Christian, E.J. and A. S. Goggi. 2008. Aromatic plant oils as fungicide for organic corn production.)



Pathogen (*Fusarium*) used in the experiment

### Leveraged funds

A \$39,672 grant was obtained from the Environmental Protection Agency, Region 7, to fund research on “Use of plant essential oils as seed treatments,” by Goggi, Rick Exner and Gary Munkvold.



Pathogen (*Penicillium*) used in the experiment

Photos by Erik J. Christian

For more information, contact Susana Goggi, 166 Seed Science Center, Iowa State University, Ames, Iowa 50011; (515) 294-6372, e-mail [susana@iastate.edu](mailto:susana@iastate.edu)